

Criteria of unconditional entanglement purification

Xiang-Bin Wang*

Imai Quantum Computation and Information project, ERATO, Japan Sci. and Tech. Agency
Daini Hongo White Bldg. 201, 5-28-3, Hongo, Bunkyo, Tokyo 113-0033, Japan

We show that the fidelity result of entanglement purification protocol with 2-way classical communications (C. H. Bennett et al, Phys. Rev. Lett., 76, 722(1996)) is always correct with whatever initial states. We then give a general criteria on the unconditional entanglement purification: if a protocol satisfies our condition, the fidelity result of the protocol with the product form of initial state for the raw pairs is also correct for the same protocol with arbitrary initial state for of the raw pairs. With this general condition, we conclude that all existing purification protocols can work unconditionally.

Quantum entanglement is one of the most important resources in quantum information processing [1]. There are purification protocols on how to distill almost perfect entangled pairs (EPR pairs) from a larger number lower quality pairs, by only local operations and classical communications. There are different types of entanglement purification protocols [2–6]. In the earliest protocols [2–4], it was assumed that those raw pairs shared by Alice and Bob are in a product state: there are no entanglement among those pairs. The idea of entanglement purification was then used to the very important problem of security proof of quantum key distribution (QKD) [8]: if Alice and Bob can share almost perfect EPR pairs through purification, they can share a secret key by measuring each pairs in the same basis on each side. This idea first appeared in Ref [4], with the advantage distillation protocol [3]. However, since the purification protocol itself assumes the product state for the initially shared raw pairs, it was concerned in Ref [4] that: “if Eve provides pairs which are entangled with each other”, then the remained pairs after distillation “may not converge to the pure state $|\phi^+\rangle\langle\phi^+|$ ”. It was then shown by Lo and Chau [7] that the result of hashing protocol given by Bennett et al [2] is actually correct with whatever form of initial states therefore a type of security proof for QKD was strictly set up. After that, Shor and Preskill show that Lo-Chau’s security proof can be reduced to the BB84 protocol which contains no difficult techniques such as quantum storage or collective measurements. (We shall call this type of protocol as prepare-and-measure protocol.) In the whole security proof of the BB84 [8] protocol based on viewpoint of entanglement purification, there are two important steps: (1), the entanglement purification protocol actually works unconditionally, with whatever type of initial state; (2), the purification can be clas-

sicalized therefore on can reduce it to a type of prepare-and-measure QKD protocol, the standard BB84. Latter, the second step was studied in a more general background and a general condition on classicalization was given [6]. With such a generalization, one does not have to specifically construct the classicalization step in a new protocol. Instead, one may directly use the main theorem in Ref [6]. However, so far there is no general statement on step 1, the unconditional entanglement purification. Say, given a purification protocol with product form of initial state for the raw pairs, under which condition the fidelity result of the final pairs is still correct given whatever initial state for the raw pairs. In this Letter, we study this problem and shall give a general condition for the unconditional entanglement purification. With our theorem, we find that all existing purification protocols can work unconditionally.

For clarity, we shall consider the details of a special case first, the (modified) advantage distillation [3,4]. This is a 2-pair error-rejection protocol with 2-way classical communications (2-CC). We then state our general theorem and consider other entanglement purification protocols [2,6] with our theorem. Note that so far no one has shown explicitly that the 2-CC protocol works unconditionally with whatever type of channel noise, though the question to protocol with 1 way classical communications (1-CC) [2] has been studied [7]. Our main contribution here is to give a general statement of the condition for unconditional entanglement purification. Since our proof is closely related to Lo-Chau’s security proof [7], we regard our result as generalized effects of Lo-Chau’s security proof.

Let’s first state our main idea. If the initial state is not a product form, classical statistics does not necessarily work. Therefore we should not blindly assume anything of classical statistics, including the error estimation from samples. To make everything clear, we shall start from a silly protocol where Alice and Bob take a dark Bell measurement (i.e. a Bell measurement without reading the measurement result) to each raw pairs. After this step, we can safely define the rates of each Bell states for the raw pairs by classical statistics. We regard state $|\phi^+\rangle$ as the perfect EPR state. *Therefore in the case that a dark Bell measurement has been done to each raw pairs, the state for raw pairs is in a product form.* In the estimation, we let Alice and Bob do collective measurements XX, YY, ZZ to test pairs. These collective measurements are just the parity measurement in different bases, e.g., the measurement result of XX is 0 or 1 if the bit values in X basis in two sides of the measured

*email: wang@qci.jst.go.jp

pair are same or different. Since all these parity measurements commute, classical sampling theory works exactly in estimating the rate of different Bell states. After measuring the test pairs, they know the distribution over 4 Bell states for the remained raw pairs by the classical sampling theory. Also, in our initial protocol, all operations commute and classical statistics works throughout the whole protocol. The fidelity between the finally distilled m pairs and the m products of perfect EPR state ($|\phi^+\rangle$) is given by classical statistics. We then show that the initial dark Bell measurement to each raw pairs can be deleted and all the collective measurements involved in the protocol can be replaced by local measurement on X, Y, Z basis in each side with the fidelity of the distilled m pairs being unchanged. Eve's information is limited by the Holevo bound [7] therefore if the fidelity of the distilled m pairs is sufficiently close to 1, her amount of information to the final key is negligible.

We denote Z, X, Y as measurement in the bases of $\{|0\rangle, |1\rangle\}$, $\{|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)\}$, $\{|y\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm i|1\rangle)\}$ respectively. We use the notation $|\phi^\pm\rangle, |\psi^\pm\rangle$ for 4 Bell states $\frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle), \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)$. We shall use WW for collective measurement to a pair in W basis and $W \otimes W$ for local measurement in W basis in each side, where W can be X, Y or Z . In the purification protocol we assume initially Alice and Bob share N raw pairs and our motivation is to distill m very good pairs finally, with its fidelity to state $(|\phi^+\rangle\langle\phi^+|)^m$ being exponentially close to 1.

Consider the following (silly) entanglement purification protocol with everything clearly defined:

Protocol 1:

Step 1: Entanglement distribution: Alice prepares N perfect EPR pairs, $|\phi^+\rangle$. She sends half of each pair to Bob (over a noisy channel in general). N is a very large natural number. Step 2: Dark Bell measurement: They take a Bell measurement to each pair but don't read the measurement results. Step 3: Error test. They randomly pick out $3k$ pairs and equally divide them into 3 groups. Note that k is large but $k \ll N$. They take collective measurement of XX to each pair in the first group, YY to each pair in the second group and ZZ to each pair in the third group. They then send the pairs in group 1,2,3 to trash can Tx, Ty, Tz , respectively. If too many of the measurement results are 1 they abort the protocol; otherwise they continue. Step 4: Error rejection. a.) Bit-flip error rejection. They randomly group the remained pairs outside the trash cans with each group containing 2 pairs. To each 2-pair group, they collect the parity information to one pair (the destination pair) by a controlled-NOT [1] operation on each side (bi-CNOT) in Z -basis. They take a collective measurement of ZZ to the destination pair. If the result is 0, they put the destination pair to the trash can Tz and keep the other pair; if it is 1, they send the destination pair to trash can Tz and discard the other pair permanently. b.) Phase-flip error rejection. They randomly group the remained pairs outside

the trash cans with each group containing 2 pairs. To each 2-pair group, they collect the X -basis parity information to the destination pair by an X -basis bi-CNOT operation. They take a collective measurement of XX to the destination pair. If the result is 0, they send the destination pair to the trash can Tx and keep the other pair, if it is 1, they send the destination to trash can Tx and discard the other pair permanently. They do operations in a.) and b.) iteratively until they believe that the state of the remained pairs outside the trash cans is very close to a product state of $|\phi^+\rangle\langle\phi^+|$. We assume there are m pairs remained outside the trash cans then.

We call this protocol as a silly protocol because it is not useful in any real entanglement purification or QKD, due to the fact that Alice and Bob have to come together to make the collective measurements. However, it is an effective mathematical tool to clearly show the security of the final useful protocol.

We regard a pair as a good pair if it is in state $|\phi^+\rangle$; a bad pair if it is in any one of the other 3 Bell states. Note that a good pair always has the parity 0 in what ever bases of X, Y, Z , while there is always a certain basis where a bad pair has the parity 1 for sure. After Step 2 all quantum entanglement among different pairs are removed therefore classical statistics works exactly with all measurement bases being commute. In particular, after Step 3, the probability of obtaining more than δN 1s by measurement WW on the remained pairs and finding more than $(\delta - \epsilon_0)k$ 1s by measurement WW on a group of k pairs is less than $\exp[-\frac{1}{4}\epsilon_0^2 N/(\delta - \delta^2)]$. Given the joint information of parity value distribution in 3 bases, one may calculate the rate of good pairs and each of the 3 types of bad pairs in remained $N - 3k$ raw pairs. We denote q_I, q_x, q_y, q_z for the rate of $|\phi^+\rangle, |\psi^+\rangle, |\psi^-\rangle, |\phi^-\rangle$. The different rates of each of the 4 Bell states after and before bit-flip error rejection are related by the following formulas [9]

$$\begin{cases} q'_I = \frac{p_I^2 + p_z^2}{(q_I + q_z)^2 + (q_x + q_y)^2}, \\ q'_x = \frac{q_x^2 + q_y^2}{(q_I + q_z)^2 + (p_x + p_y)^2}, \\ q'_y = \frac{2q_x q_y}{(q_I + q_z)^2 + (q_x + q_y)^2}, \\ q'_z = \frac{2q_I q_z}{(q_I + q_z)^2 + (q_x + p_y)^2}. \end{cases} \quad (1)$$

Here the left hand sides are the quantities after bit-flip rejection operation and the symbols in right hand sides of each equations are for the quantities before bit-flip rejection operation. Switching q_x and q_z we shall obtain the relations after and before phase-flip rejection operation. This shows that part a.) of Step 4 reduces the rate of bad pairs of $|\psi^\pm\rangle$; part b.) reduces the rate of bad

pairs of $|\phi^-\rangle, |\psi^-\rangle$. Also, it can be shown that, the rate of remained pairs after the Bit-flip rejection is

$$f = \frac{1}{2(q_I + q_z)^2 + (q_x + p_y)^2}. \quad (2)$$

If the initial error rates are small (say, $q_x = q_y = q_z = t_0$), after g rounds of error rejection, the rate of remained pairs is in the magnitude order of 4^{-g} while the new bad-pair rate is in the magnitude order of $4^g t_0^{2g}$. If t_0 is small, a few steps of repetition of error-rejection is sufficient. This has been demonstrated numerically by Deutsch et al [4]. We conclude that with an appropriate number of iteration, the state of the finally remained m pairs ρ_m must be exponentially close to m perfect EPR pairs, i.e.

$$\langle \Phi_m | \rho_m | \Phi_m \rangle = 1 - \epsilon \quad (3)$$

and ϵ is a small number exponentially close to 0, $|\Phi_m\rangle = |\phi^+\rangle^{\otimes m}$.

The order of Step 2 and Step 3 can be exchanged since the collective measurements in Step 3 are just coarse-grained Bell measurements which commute with the dark Bell measurements in Step 2. For example, measurement ZZ is simply a measurement on whether the pair belongs to subspace $\{|\phi^\pm\rangle\}$ or subspace $\{|\psi^\pm\rangle\}$.

Consider the bi-CNOT in Z basis. On each side they take the unitary operation of $|0\rangle|0\rangle \rightarrow |0\rangle|0\rangle, |0\rangle|1\rangle \rightarrow |0\rangle|1\rangle, |1\rangle|0\rangle \rightarrow |1\rangle|1\rangle, |1\rangle|1\rangle \rightarrow |1\rangle|0\rangle$, with the second state being the destination qubit. Consequently, if the second pair is destination pair, a Z -basis bi-CNOT operation is a permutation in Bell basis as the following

$$|\chi_{i,j}\rangle|\chi_{i',j'}\rangle \longrightarrow |\chi_{i\oplus i',j}\rangle|\chi_{i',j'\oplus j}\rangle. \quad (4)$$

Here i, j can be either 0 or 1, $|\chi_{0,0}\rangle, |\chi_{1,0}\rangle, |\chi_{0,1}\rangle, |\chi_{1,1}\rangle$ are notations for $|\phi^+\rangle, |\phi^-\rangle, |\psi^+\rangle, |\psi^-\rangle$, respectively and symbol $\oplus$ is the calculation of summation mod 2. This permutation operation and the dark Bell measurement commute. *Proof:* Given arbitrary l -pair state ρ_l , one can always write in the form $\rho_l = \sum_l |\langle b_l | \rho_l | b_l \rangle \cdot b_l \rangle \langle b_l| + \hat{O}_{off}$, where $\{|b_l\rangle\}$ is the Bell bases for l pairs, l is from 1 to 4^l , $\hat{O}_{off}$ contains all off-diagonal terms in Bell bases which have no contribution to the outcome of Bell measurement. With the original order of the protocol, i.e., first take dark measurement and then take bi-CNOTs, the outcome state is $\rho_f = \hat{P}(\sum_l |\langle b_l | \rho_l | b_l \rangle \cdot b_l \rangle \langle b_l|)$ and $\hat{P}$ is the permutation given by bi-CNOTs. If we reverse the order, after the bi-CNOTs, the state is changed to $\hat{P}(\sum_l |\langle b_l | \rho_l | b_l \rangle \cdot b_l \rangle \langle b_l|) + \hat{P}(\rho_{off})$. Note that with the permutation $\hat{P}$ in Bell bases, the off-diagonal terms will be still off-diagonal and the diagonal terms will be still diagonal. If we then take the Dark Bell measurement, only the diagonal terms survive. This is to say, with the order being reversed, the outcome is also ρ_f . This makes the proof.

Similarly, one can show that the X -basis bi-CNOT also commutes with the dark Bell measurement. Therefore

all bi-CNOT operations commute with the Bell measurement in Step 2. Therefore all operations in Step 4 and Step 2 commute, consequently we can exchange the order of Step 2 and Step 4, i.e. Step 2 in protocol 1 can be delayed until the end of the protocol. This dark Bell measurement in the end of the protocol can be deleted because it does not change the fidelity of the remained m pairs. Explicitly, given any state of l pairs, ρ_l , after a dark Bell measurement to each pair, the state is changed to $\rho'_l = \sum_i \langle b_i | \rho'_l | b_i \rangle \cdot |b_i\rangle \langle b_i|$, with $\{|b_i\rangle\}$ being l -Bell basis and i running from 1 to 4^l . The fidelity of ρ'_l is

$$\langle b_1 | \rho'_l | b_1 \rangle = \langle b_1 | \sum_i \langle b_i | \rho'_l | b_i \rangle \cdot |b_i\rangle \langle b_i| | b_1 \rangle = \langle b_1 | \rho_l | b_1 \rangle \quad (5)$$

and $|b_1\rangle = |\phi^+\rangle^{\otimes l}$. This shows that, although the dark Bell measurement may change the state of remained m pairs, it does not change the fidelity. Due to the above arguments, we now give the following modified protocol for QKD:

Protocol 2:

Step 1: Entanglement distribution. Step 2: Error test. Step 3: Error rejection. Step 4: Alice and Bob send all trash cans to Eve. Step 5: They measure $Z \otimes Z$ to each of the remained m pairs outside the trash cans and use the result as an m -bit final key.

Step 1-3 are same with those in Protocol 1. As it has been argued already, after Step 3 in protocol 2, the state of remained m pairs is exponentially close to 1. Therefore Eve's information to the final key is upper bounded by a negligibly small number ϵ' [7], even Eve controls everything except for the distilled m pairs, including all pairs in trash cans and everything else outside Alice and Bob's labs.

This protocol is still useless for any actual entanglement purification because it requires the collective measurements though the step of dark Bell measurement to each pair has now been removed. Now we show that if we replace all collective measurements WW by local measurements $W \otimes W$, the final result of the protocol is unchanged.

For such a purpose, we first show that Step 4 can be replaced by Step 4':

Step 4': Alice and Bob don't send any states in trash cans to Eve. Instead they take local measurement of $W \otimes W$ to the pairs in trash cans Tw , (Tw can be Tx , Ty or Tz .) and announce the results.

If we only want an unconditionally secure final key, we can use the following simple argument: A restrictive Eve cannot do the attack better than an un-restricted Eve. Eve in protocol 2 is an un-restricted Eve who controls everything except for the distilled m pairs. Consider a restriction requires Eve to measure $W \otimes W$ to each pairs in trash cans Tw and announce the measurement results. Note that except for this, Eve's action to everything else under her control is not restricted, e.g., the ancillas which had been prepared in the most initial stage when Alice distributes EPR pairs to Bob. With

the above restriction, Eve's information to the final key must be also bounded by ϵ' . Or in other words, if Eve's information is bounded by ϵ' with *whatever* attack, her information must be also bounded by ϵ' with any *specific* type of attack. Replacing Step 4 with Step 4' in protocol 2 is equivalent to add the above restriction to Eve in protocol 2. This is to say, it will cause no harm to the security of final key if Alice and Bob keep all trash cans and take local measurement $W \otimes W$ to pairs in trash cans Tw and announce the measurement outcome.

We can also show that Step 4 can be replaced by Step 4' by the fidelity criteria, from the viewpoint of entanglement purification. Consider the case that Step 4 is replaced Step 4'. We only need to argue that the local measurement in Step 4' may cause no changes to the fidelity of the m remained pairs. Suppose the state of the total space immediately after Step 3 is $\rho_t = \rho_{m,c,e}$ and the subscript m, c, e denote for the subspace of m remained pairs, trash cans and environment, respectively. The state of the remained m pairs immediately after Step 3 is

$$\rho_m = tr_{c,e} \rho_t. \quad (6)$$

The notation $tr_{c,e}$ means the partial trace calculation in subspaces of trash cans and environment. In the protocol, if the error test is passed, Alice and Bob will *always* accept the remained m pairs no matter what the measurement outcome they may obtain in Step 4'. Therefore for the quantity of fidelity of the remained m pairs, we only care about the *averaged* state for the m pairs after Step 4', with the average being taken over all possible measurement results in Step 4'. Mathematically, the averaged total state after Step 4' is

$$\rho'_t = \sum_L \langle L | \rho_t | L \rangle | L \rangle \langle L|. \quad (7)$$

Here $|L\rangle$ is an eigenstate of the local measurement to those qubits in trash cans in Step 4'. Therefore the averaged state for the remained m pairs after Step 4' is

$$\rho'_m = tr_{c,e} \rho'_t = tr_e \sum_L \langle L | \rho_t | L \rangle = tr_{c,e} \rho_t = \rho_m. \quad (8)$$

This shows that the averaged state for the remained m pairs after Step 4' is just ρ_m , therefore Step 4' does not change the fidelity of the remained m pairs.

Now we show why all collective measurements WW can simply be replaced by local measurements $W \otimes W$. With Step 4 being replaced by Step 4', each pairs in trash cans are measured two times, the first is WW , the second is $W \otimes W$. They commute, therefore one can exchange the order of them. More over, once we have measured $W \otimes W$, we do not need to measure WW , since the result of WW has been explicitly determined already by the measurement $W \otimes W$. Therefore all the collective measurements in protocol 2 can be replaced by local measurements in each side. We obtain the following protocol

with all measurements being local:

Protocol 3:

Step 1: Entanglement distribution. Step 2: Error test. They randomly pick out $3k$ pairs and equally divide them into 3 groups. They take local measurement of $X \otimes X, Y \otimes Y, Z \otimes Z$ to each pair in group 1,2,3, respectively. They announce the measurement results and discard those measured qubits. If too many of the results from two sides disagree, they abort the protocol; otherwise they continue. Step 3:a.) Bit-flip error rejection. They randomly group the remained pairs with each group containing 2 pairs. To each 2-pair group, they collect the parity information to the destination pair by a Z -basis bi-CNOT operation. They take local measurement $Z \otimes Z$ to the destination pair and announce the measurement results from two sides. If they are same, they discard the destination pair and keep the control pair; if they are different they discard both pairs. b.) Phase-flip error rejection. They randomly group the remained pairs with each group containing 2 pairs. To each 2-pair group, they collect the X -basis parity information to the destination pair by a X -basis bi-CNOT operation. They measure $X \otimes X$ to the destination pair and announce the measurement results from two sides. If they are same, they discard the destination pair and keep the control pair; if they are different they discard both pairs. They do operations in a.) and b.) iteratively and finally there are m pairs remained outside the trash cans. Step 4: They measure $Z \otimes Z$ to each of the remained m pairs and use the result as an m -bit final key.

So far we have given a detailed and rigorous proof of the entanglement purification and security of QKD with 2-CC. There are two important issues in the reduction. The first issue is the removal of the initial dark Bell measurement: this measurement commute with bi-CNOTs and all collective measurements of WW , therefore it can be delayed and removed, with the fidelity of the purification outcome unchanged. The second issue is, after removal of dark Bell measurement, any collective measurements WW can be replaced by local measurement of W on each side, i.e., $W \otimes W$, since WW and $W \otimes W$ commute. Therefore we propose the following general theorem:

Theorem: *Suppose we have certain fidelity result for the final pairs through a certain purification protocol with product form of state for the raw pairs. We must have the same fidelity result for the same protocol with whatever form of state for the raw pairs, provided that the protocol satisfies the following conditions: (1) All quantum operations needed are not more than bi-CNOTs and local measurements on each side, i.e. $\{W \otimes W\}$; (2) The fidelity result is unchanged if we replace any local measurement $W \otimes W$ by collective measurement WW ; (3) All WW commute with dark Bell measurement.*

Together with the main theorem in Ref [6], this theorem works effectively on the security proof of any purification-based QKD protocol. We can just consider the conditional purification first: the special case of prod-

uct state for the raw pairs and consider the fidelity result of purification. After that, we use our theorem and the main theorem in Ref [6] to reduce it to the unconditional entanglement purification and moreover, prepare-and-measure QKD protocol.

To reduce protocol 3 to a prepare-and-measure protocol, we only need to replace the Phase-flip rejection part in Step 3 with 3-pair phase error correction used in Ref [6]. There are only bi-CNOTs and collective measurements $X \otimes X, Z \otimes Z$ involved [6]. According to our theorem, the purification must be also unconditional. Therefore the unconditional security of QKD [6,9,10] protocols with 2-CC is confirmed. Similarly, the entanglement purification protocol with hashing [2] and the one with quantum error correction code [5] also only contains bi-CNOTs and local measurement $W \otimes W$ therefore the fidelity result of the protocol is unconditionally true with whatever initial state. The issue of side information of Shor-Preskill protocol has been studied by Hwang [11].

Acknowledgement: I thank Prof Imai H for support. I thank H. P Yuen for comments.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, 2000.
 - [2] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A54, 3824(1996).
 - [3] C. H. Bennett, G. Brassard, S. Popescu, B. Schmachter, J. Smolin, and W. K. Wootters, Phys. Rev. Lett., 76, 722(1996).
 - [4] D. Deutsch, A. Ekert, R. Jozsa, C. Macchiavello, S. Popescu, and A. Sanpera, Phys. Rev. Lett., 77, 2818(1996); Erratum Phys. Rev. Lett. **80**, 2022 (1998).
 - [5] P. W. Shor and J. Preskill, Phys. Rev. Lett., vol. 85,441(2000).
 - [6] D. Gottesman and H.-K. Lo, IEEE Transactions on Information Theory, 49, 457(2003).
 - [7] H.-K. Lo and H. F. Chau, Science, 283, 2050(1999).
 - [8] C. H. Bennett and G. Brassard, *Proceedings of IEEE International Conference on Computers, Systems and Signal Processing, Bangalore, India, 1984*, (IEEE Press, 1984), pp. 175–179; C.H. Bennett and G. Brassard, IBM Technical Disclosure Bulletin **28**, 3153–3163 (1985).
 - [9] H. F. Chau, Phys. Rev. A66, 060302(R) (2002).
 - [10] X. B. Wang, Phys. Rev. Lett., 92, 077902(2004).
 - [11] W. Y. Hwang, quant-ph/0402032.